

HAIR ELEMENTS REPORT INTRODUCTION

Hair is an excretory tissue for essential, nonessential and potentially toxic elements. In general, the amount of an element that is irreversibly incorporated into growing hair is proportional to the level of the element in other body tissues. Therefore, hair elements analysis provides an indirect screening test for physiological excess, deficiency or maldistribution of elements in the body. Clinical research indicates that hair levels of specific elements, particularly potentially toxic elements such as cadmium, mercury, lead and arsenic, are highly correlated with pathological disorders. For such elements, levels in hair may be more indicative of body stores than the levels in blood and urine.

All screening tests have limitations that must be taken into consideration. The correlation between hair element levels and physiological disorders is determined by numerous factors. Individual variability and compensatory mechanisms are major factors that affect the relationship between the distribution of elements in hair and symptoms and pathological conditions. It is also very important to keep in mind that scalp hair is vulnerable to external contamination of elements by exposure to hair treatments and products. Likewise, some hair treatments (e.g. permanent solutions, dyes, and bleach) can strip hair of endogenously acquired elements and result in false low values. Careful consideration of the limitations must be made in the interpretation of results of hair analysis. The data provided should be considered in conjunction with symptomology, diet analysis, occupation and lifestyle, physical examination and the results of other analytical laboratory tests.

Caution: The contents of this report are not intended to be diagnostic and the physician using this information is cautioned against treatment based solely on the results of this screening test. For example, copper supplementation based upon a result of low hair copper is contraindicated in patients afflicted with Wilson's Disease.

Aluminum High

The Aluminum (Al) level in hair is a reliable indicator of assimilation of this element, provided that hair preparations have not added exogenous Al. Al is a nonessential element that can be toxic if excessively assimilated into cells.

Excess Al can inhibit the formation of alpha-keto glutarate and result in toxic levels of ammonia in tissues. Al can bond to phosphorylated bases on DNA and disrupt protein synthesis and catabolism. Al excess should be considered when symptoms of presenile dementia or Alzheimer's disease are observed. Hair Al is commonly elevated in children and adults with low zinc and behavioral/learning disorders such as ADD, ADHD and autism. Individuals with renal problems or on renal dialysis may have elevated Al.

Possible sources of Al include some antacid medications, Al cookware, baking powder, processed cheese, drinking water, and antiperspirant components that may be absorbed. Analyses performed at DDI indicate extremely high levels of Al are in many colloidal mineral products.

Al has neurotoxic effects at high levels, but low levels of accumulation may not elicit immediate symptoms. Early symptoms of Al burden may include: fatigue, headache, and symptoms of phosphate depletion.

A urine elements test can be used to corroborate Al exposure. Al can be effectively complexed and excreted with silicon (J. Environ. Pathol. Toxicol. Oncol., 13(3): 205-7, 1994). A complex of malic acid and Mg has been reported to be quite effective in lowering Al levels (DDI clients).

Antimony High

Hair is a preferred tissue for analysis of Antimony (Sb) exposure and body burden. Elevated hair Sb levels have been noted as long as a year after exposure.

Sb is a nonessential element that is chemically similar to but less toxic than arsenic. Food and smoking are the usual sources of Sb. Thus cigarette smoke can externally contaminate hair, as well as contribute to uptake via inhalation. Gunpowder (ammunition) often contains Sb. Firearm enthusiasts often have elevated levels of Sb in hair. Other possible sources are textile industry, metal alloys, and some antihelminthic and antiprotozoic drugs. Sb is also used in the manufacture of paints, glass, ceramics, solder, batteries, bearing metals and semiconductors.

Like arsenic, Sb has a high affinity for sulfhydryl groups on many enzymes. Sb is conjugated with glutathione and excreted in urine and feces. Therefore, excessive exposure to Sb has the potential to deplete intracellular glutathione pools.

Early signs of Sb excess include: fatigue, muscle weakness, myopathy, nausea, low back pain, headache, and metallic taste. Later symptoms include hemolytic anemia, myoglobinuria, hematuria and renal failure. Transdermal absorption can lead to "antimony spots" which resemble chicken pox. Respiratory tissue irritation may result from inhalation of Sb particles or dust.

A confirmatory test for recent or current exposure is the measurement of Sb in the urine.

Cadmium high

Hair Cadmium (Cd) levels provide an excellent indication of body burden. Cd is a toxic heavy metal that has no metabolic function in the body. Moderately high Cd levels, about 4-8 µg/g, may be associated with hypertension, while very severe Cd toxicity may cause hypotension. Cd adversely affects the kidneys, lungs, testes, arterial walls, and bones and interferes with many enzymatic reactions. Chronic Cd excess can lead to microcytic, hypochromic anemia and proteinuria with loss of beta-2-microglobulin, and functional zinc deficiency. Cd excess is also commonly associated with fatigue, weight loss, osteomalacia, and lumbar pain.

Cd absorption is reduced by zinc, calcium, and selenium. Cd is found in varying amounts in foods, from .04 µg/g for some fruits to 3-5 µg/g in some oysters and anchovies. Cigarette smoking significantly increases Cd intake. Refined carbohydrates have very little zinc in relation to the Cd.

If hair zinc is not abnormal, external contamination may have caused the elevated hair Cd level. Exogenous contamination may come from permanent solutions, dyes, bleach, and some hair sprays. A confirming test for elevated body burden of Cd is urine analysis following administration of appropriate chelating agents: EDTA, sulfhydryl agents (DMSA, D-penicillamine, DMPS).

Potassium High

High hair Potassium (K) is not necessarily reflective of dietary intake or nutrient status. However, elevated K may be reflective of metabolic disorders associated with exposure to potentially toxic elements.

K is an electrolyte and a potentiator of enzyme functions, but neither of these functions take place in hair. Elevated K in hair may reflect overall retention of K by the body or maldistribution of this element. In adrenocortical insufficiency, K is increased in blood, while it is decreased in urine; cellular K may or may not be increased. Also, hair is occasionally contaminated with K from some shampoos. Observations at DDI indicate that K and sodium levels in hair are commonly high in association with toxic element burden. The elevated K and sodium levels are often concomitant with low levels of calcium and magnesium in hair. This apparent phenomena requires further investigation.

Elevated hair potassium should be viewed as a screening test. Appropriate tests for excess body K include measurements of packed red blood cell K; serum or whole blood K and sodium/K ratio, measurement of urine K and sodium/K ratio; and an assessment of adrenocortical function.

Copper High

The high level of Copper (Cu) in hair may be indicative of excess Cu in the body. However, it is important first to rule out exogenous contamination sources: permanent solutions, dyes, bleaches, swimming pool/hot tub water, and washing hair in acidic water carried through Cu pipes. In the case of contamination from hair preparations, other elements (aluminum, silver, nickel, titanium) are usually also elevated.

Sources of excessive Cu include contaminated food or drinking water, excessive Cu supplementation, and occupational or environmental exposures. Insufficient intake of competitively absorbed elements such as zinc or molybdenum can lead to, or worsen Cu excess.

Medical conditions that may be associated with excess Cu include: biliary obstruction (reduced ability to excrete Cu), liver disease (hepatitis or cirrhosis), and renal dysfunction. Symptoms associated with excess Cu accumulation are muscle and joint pain, depression, irritability, tremor, hemolytic anemia, learning disabilities, and behavioral disorders.

Confirmatory tests for Cu excess are a comparison of Cu in pre vs. post provocation (D-penicillamine, DMPS) urine elements tests and a whole blood elements analysis.

Vanadium High

High levels of Vanadium (V) in hair may be indicative of excess absorption of the element. It is well established that excess V can have toxic effects in humans. Although it appears that V may have essential functions, over zealous supplementation is not warranted.

Excess levels of V in the body can result from chronic consumption of fish, shrimp, crabs, and oysters derived from water near offshore oil rigs (Metals in Clinical and Analytical Chemistry, 1994). Industrial/environmental sources of V include: processing of mineral ores, phosphate fertilizers, combustion of oil and coal, production of steel, and chemicals used in the fixation of

dyes and print.

Symptoms of V toxicity vary with chemical form and route of absorption. Inhalation of excess V may produce respiratory irritation and bronchitis. Excess ingestion of V can result in decreased appetite, depressed growth, diarrhea/gastrointestinal disturbances, nephrotoxic and hematotoxic effects. Pallor, diarrhea, and green tongue are early signs of excess V and have been reported in human subjects consuming about 20 mg V/day (Modern Nutrition in Health and Disease, 8th edition, eds. Shils, M., Olson, J., and Mosha, S., 1994).

Confirmatory tests for excess V are red blood cell elements analysis, and urine V which reflects recent intake.

Selenium Low

Selenium (Se) is normally found in hair at very low levels, and several studies provide evidence that low hair Se is reflective of dietary intake and associated with cardiovascular disorders. Utilization of hair Se levels to assess nutritional status, however, is complicated by the fact that use of Se- or sulfur-containing shampoo markedly increases hair Se (externally) and can give a false high value.

Se is an extremely important essential element due to its antioxidative function as an obligatory component of the enzyme glutathione peroxidase. Se is also protective in its capacity to bind and "inactivate" mercury, and Se is an essential cofactor in the deiodination of T-4 to active T-3 (thyroid hormone). Some conditions of functional hypothyroidism therefore may be due to Se deficiency (Nature; 349:438-440, 1991); this is of particular concern with mercury exposure. Studies have also indicated significant inverse correlations between Se and heart disease, cancer, and asthma.

Selenium deficiency is common and can result from low dietary intake of Se or vitamin E, and exposure to toxic metals, pesticides/herbicides and chemical solvents.

Symptoms of Se deficiency are similar to that of vitamin E deficiency and include muscle aches, increased inflammatory response, loss of body weight, alopecia, listlessness, skeletal and muscular degeneration, growth stunting, and depressed immune function.

Confirmatory tests for Se deficiency are Se content of packed red blood cells, and activity of glutathione peroxidase in red blood cells.

Strontium High

Hair usually reflects the body burden of Strontium (Sr), and Sr levels usually correlate with calcium levels in body tissue. However, hair levels of Sr can be raised by external contamination, usually from hair treatment products. Elevated Sr in hair treated with permanent solutions, dyes, or bleaches is likely to be an artifact of hair treatment and probably does not reflect the level of Sr in other tissues.

Diseases of excess Sr have not been reported, except for Sr rickets. In general, Sr excess is not of clinical concern in the U.S. It's bad reputation comes from it's radioactive isotopes which were widespread in the western U.S. as a result of nuclear testing in the 1950's. Stable Sr (not

radioactive Sr) is measured and reported by DDI.

Other tests indicative of Sr status or excess are measurements of Sr in whole blood, Sr/calcium ratio in blood, and Sr in urine.

Total Toxic Element Indication

The potentially toxic elements vary considerably with respect to their relative toxicities. The accumulation of more than one of the most toxic elements may have synergistic adverse effects, even if the level of each individual element is not strikingly high. Therefore, we present a total toxic element "score" which is estimated using a weighted average based upon relative toxicity. For example, the combined presence of lead and mercury will give a higher total score than that of the combination of silver and beryllium.

Lab number: **H060327-0290-1**
Patient: **Harley Lunn**

Hair

Page: 8
Client: **29327**
